

## GATE 2022 Mathematics (MA)

### Useful data

$A \setminus B$	$\{a \in A : a \notin B\}$
$\mathbb{C}$	Set of all complex numbers
$\mathbb{C}^{m \times n}$	Set of all matrices of order $m \times n$ with complex entries
$\mathbb{C}^\infty(\Omega)$	Collection of all infinitely differentiable functions on the open domain $\Omega$
$i$	$\sqrt{-1}$
$I$	Identity matrix of appropriate order
$L^2(\mathbb{R})$	$:= L^2(\mathbb{R}, dx)$
$L^2[a, b]$	$:= L^2([a, b], dx)$
$\mathbb{N}$	Set of all positive integers
$\mathbb{Q}$	Set of all rational numbers
$\mathbb{R}$	Set of all real numbers
$\mathbb{R}^{m \times n}$	Set of all matrices of order $m \times n$ with real entries
$\mathbb{S}^1$	$\{(x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 = 1\}$
$\mathbb{S}^2$	$\{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1^2 + x_2^2 + x_3^2 = 1\}$
$\mathbb{Z}$	Set of all integers



**GATE 2022 General Aptitude (GA)**

**Q.1 – Q.5 Carry ONE mark each.**

Q.1	As you grow older, an injury to your _____ may take longer to _____ .
(A)	heel / heel
(B)	heal / heel
(C)	heal / heal
(D)	heel / heal




Q.2	<p>In a 500 m race, P and Q have speeds in the ratio of 3 : 4. Q starts the race when P has already covered 140 m.</p> <p>What is the distance between P and Q (in m) when P wins the race?</p>
(A)	20
(B)	40
(C)	60
(D)	140



Q.3	<p>Three bells P, Q, and R are rung periodically in a school. P is rung every 20 minutes; Q is rung every 30 minutes and R is rung every 50 minutes.</p> <p>If all the three bells are rung at 12:00 PM, when will the three bells ring together again the next time?</p>
(A)	5:00 PM
(B)	5:30 PM
(C)	6:00 PM
(D)	6:30 PM



<p>Q.4</p>	<p>Given below are two statements and four conclusions drawn based on the statements.</p> <p>Statement 1: Some bottles are cups.</p> <p>Statement 2: All cups are knives.</p> <p>Conclusion I: Some bottles are knives.</p> <p>Conclusion II: Some knives are cups.</p> <p>Conclusion III: All cups are bottles.</p> <p>Conclusion IV: All knives are cups.</p> <p>Which one of the following options can be logically inferred?</p>
<p>(A)</p>	<p>Only conclusion I and conclusion II are correct</p>
<p>(B)</p>	<p>Only conclusion II and conclusion III are correct</p>
<p>(C)</p>	<p>Only conclusion II and conclusion IV are correct</p>
<p>(D)</p>	<p>Only conclusion III and conclusion IV are correct</p>

<p>Q.5</p>	<p>The figure below shows the front and rear view of a disc, which is shaded with identical patterns. The disc is flipped once with respect to any one of the fixed axes 1-1, 2-2 or 3-3 chosen uniformly at random.</p> <p>What is the probability that the disc <b>DOES NOT</b> retain the same front and rear views after the flipping operation?</p> <div style="text-align: center;">  </div> <p style="text-align: center;">Front View                      Rear View</p>
<p>(A) 0</p>	
<p>(B) <math>\frac{1}{3}</math></p>	
<p>(C) <math>\frac{2}{3}</math></p>	
<p>(D) 1</p>	



**Q. 6 – Q. 10 Carry TWO marks each.**

<p>Q.6</p>	<p>Altruism is the human concern for the wellbeing of others. Altruism has been shown to be motivated more by social bonding, familiarity and identification of belongingness to a group. The notion that altruism may be attributed to empathy or guilt has now been rejected.</p> <p>Which one of the following is the CORRECT logical inference based on the information in the above passage?</p>
<p>(A)</p>	<p>Humans engage in altruism due to guilt but not empathy</p>
<p>(B)</p>	<p>Humans engage in altruism due to empathy but not guilt</p>
<p>(C)</p>	<p>Humans engage in altruism due to group identification but not empathy</p>
<p>(D)</p>	<p>Humans engage in altruism due to empathy but not familiarity</p>



<p>Q.7</p>	<p>There are two identical dice with a single letter on each of the faces. The following six letters: Q, R, S, T, U, and V, one on each of the faces. Any of the six outcomes are equally likely.</p> <p>The two dice are thrown once independently at random.</p> <p>What is the probability that the outcomes on the dice were composed only of any combination of the following possible outcomes: Q, U and V?</p>
<p>(A)</p>	<p><math>\frac{1}{4}</math></p>
<p>(B)</p>	<p><math>\frac{3}{4}</math></p>
<p>(C)</p>	<p><math>\frac{1}{6}</math></p>
<p>(D)</p>	<p><math>\frac{5}{36}</math></p>










Q.8	<p>The price of an item is 10% cheaper in an online store S compared to the price at another online store M. Store S charges ₹ 150 for delivery. There are no delivery charges for orders from the store M. A person bought the item from the store S and saved ₹ 100.</p> <p>What is the price of the item at the online store S (in ₹) if there are no other charges than what is described above?</p>
(A)	2500
(B)	2250
(C)	1750
(D)	1500



<p>Q.9</p>	<p>The letters P, Q, R, S, T and U are to be placed one per vertex on a regular convex hexagon, but not necessarily in the same order.</p> <p>Consider the following statements:</p> <ul style="list-style-type: none"> <li>• The line segment joining R and S is longer than the line segment joining P and Q.</li> <li>• The line segment joining R and S is perpendicular to the line segment joining P and Q.</li> <li>• The line segment joining R and U is parallel to the line segment joining T and Q.</li> </ul> <p>Based on the above statements, which one of the following options is CORRECT?</p>
<p>(A)</p>	<p>The line segment joining R and T is parallel to the line segment joining Q and S</p>
<p>(B)</p>	<p>The line segment joining T and Q is parallel to the line joining P and U</p>
<p>(C)</p>	<p>The line segment joining R and P is perpendicular to the line segment joining U and Q</p>
<p>(D)</p>	<p>The line segment joining Q and S is perpendicular to the line segment joining R and P</p>



<p>Q.10</p>	<div style="text-align: center;">  </div> <p>An ant is at the bottom-left corner of a grid (point P) as shown above. It aims to move to the top-right corner of the grid. The ant moves only along the lines marked in the grid such that the current distance to the top-right corner strictly decreases.</p> <p>Which one of the following is a part of a possible trajectory of the ant during the movement?</p>
<p>(A)</p>	
<p>(B)</p>	
<p>(C)</p>	
<p>(D)</p>	

**GATE 2022 Mathematics (MA)**

**Q.11 – Q.35 Carry ONE mark each.**

Q.11	<p>Suppose that the characteristic equation of <math>M \in \mathbb{C}^{3 \times 3}</math> is</p> $\lambda^3 + \alpha\lambda^2 + \beta\lambda - 1 = 0,$ <p>where <math>\alpha, \beta \in \mathbb{C}</math> with <math>\alpha + \beta \neq 0</math>. Which of the following statements is TRUE?</p>
(A)	$M(I - \beta M) = M^{-1}(M + \alpha I)$
(B)	$M(I + \beta M) = M^{-1}(M - \alpha I)$
(C)	$M^{-1}(M^{-1} + \beta I) = M - \alpha I$
(D)	$M^{-1}(M^{-1} - \beta I) = M + \alpha I$



**GATE 2022 Mathematics (MA)**

<p>Q.12</p>	<p>Consider</p> <p><b>P:</b> Let <math>M \in \mathbb{R}^{m \times n}</math> with <math>m &gt; n \geq 2</math>. If <math>\text{rank}(M) = n</math>, then the system of linear equations <math>Mx = 0</math> has <math>x = 0</math> as the only solution.</p> <p><b>Q:</b> Let <math>E \in \mathbb{R}^{n \times n}, n \geq 2</math> be a non-zero matrix such that <math>E^3 = 0</math>. Then <math>I + E^2</math> is a singular matrix.</p> <p>Which of the following statements is TRUE?</p>
<p>(A)</p>	<p>Both <b>P</b> and <b>Q</b> are TRUE</p>
<p>(B)</p>	<p>Both <b>P</b> and <b>Q</b> are FALSE</p>
<p>(C)</p>	<p><b>P</b> is TRUE and <b>Q</b> is FALSE</p>
<p>(D)</p>	<p><b>P</b> is FALSE and <b>Q</b> is TRUE</p>

**GATE 2022 Mathematics (MA)**

Q.13	<p>Consider the real function of two real variables given by</p> $u(x, y) = e^{2x} [\sin 3x \cos 2y \cosh 3y - \cos 3x \sin 2y \sinh 3y].$ <p>Let <math>v(x, y)</math> be the harmonic conjugate of <math>u(x, y)</math> such that <math>v(0, 0) = 2</math>. Let <math>z = x + iy</math> and <math>f(z) = u(x, y) + iv(x, y)</math>, then the value of <math>4 + 2if(i\pi)</math> is</p>
(A)	$e^{3\pi} + e^{-3\pi}$
(B)	$e^{3\pi} - e^{-3\pi}$
(C)	$-e^{3\pi} + e^{-3\pi}$
(D)	$-e^{3\pi} - e^{-3\pi}$



GATE 2022 Mathematics (MA)

Q.14	<p>The value of the integral</p> $\int_C \frac{z^{100}}{z^{101} + 1} dz$ <p>where <math>C</math> is the circle of radius 2 centred at the origin taken in the anti-clockwise direction is</p>
(A)	$-2\pi i$
(B)	$2\pi$
(C)	0
(D)	$2\pi i$

**GATE 2022 Mathematics (MA)**

Q.15	Let $X$ be a real normed linear space. Let $X_0 = \{x \in X : \ x\  = 1\}$ . If $X_0$ contains two distinct points $x$ and $y$ and the line segment joining them, then, which of the following statements is TRUE?
(A)	$\ x + y\  = \ x\  + \ y\ $ and $x, y$ are linearly independent
(B)	$\ x + y\  = \ x\  + \ y\ $ and $x, y$ are linearly dependent
(C)	$\ x + y\ ^2 = \ x\ ^2 + \ y\ ^2$ and $x, y$ are linearly independent
(D)	$\ x + y\  = 2\ x\ \ y\ $ and $x, y$ are linearly dependent





GATE 2022 Mathematics (MA)

Q.16	<p>Let <math>\{e_k : k \in \mathbb{N}\}</math> be an orthonormal basis for a Hilbert space <math>H</math>.          Define <math>f_k = e_k + e_{k+1}, k \in \mathbb{N}</math> and <math>g_j = \sum_{n=1}^j (-1)^{n+1} e_n, j \in \mathbb{N}</math>.          Then <math>\sum_{k=1}^{\infty}  \langle g_j, f_k \rangle ^2 =</math></p>
(A)	0
(B)	$j^2$
(C)	$4j^2$
(D)	1



**GATE 2022 Mathematics (MA)**

Q.17	Consider $\mathbb{R}^2$ with the usual metric. Let $A = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\}$ and $B = \{(x, y) \in \mathbb{R}^2 : (x-2)^2 + y^2 \leq 1\}$ . Let $M = A \cup B$ and $N = \text{interior}(A) \cup \text{interior}(B)$ . Then, which of the following statements is TRUE?
(A)	$M$ and $N$ are connected
(B)	Neither $M$ nor $N$ is connected
(C)	$M$ is connected and $N$ is not connected
(D)	$M$ is not connected and $N$ is connected



GATE 2022 Mathematics (MA)

Q.18	<p>The real sequence generated by the iterative scheme</p> $x_n = \frac{x_{n-1}}{2} + \frac{1}{x_{n-1}}, \quad n \geq 1$
(A)	converges to $\sqrt{2}$ , for all $x_0 \in \mathbb{R} \setminus \{0\}$
(B)	converges to $\sqrt{2}$ , whenever $x_0 > \sqrt{\frac{2}{3}}$
(C)	converges to $\sqrt{2}$ , whenever $x_0 \in (-1, 1) \setminus \{0\}$
(D)	diverges for any $x_0 \neq 0$



GATE 2022 Mathematics (MA)

Q.19	The initial value problem $\frac{dy}{dx} = \cos(xy), \quad x \in \mathbb{R}, \quad y(0) = y_0,$ where $y_0$ is a real constant, has
(A)	a unique solution
(B)	exactly two solutions
(C)	infinitely many solutions
(D)	no solution



GATE 2022 Mathematics (MA)

Q.20	<p>If eigenfunctions corresponding to distinct eigenvalues <math>\lambda</math> of the Sturm-Liouville problem</p> $\frac{d^2y}{dx^2} - 3\frac{dy}{dx} = \lambda y, \quad 0 < x < \pi,$ $y(0) = y(\pi) = 0$ <p>are orthogonal with respect to the weight function <math>w(x)</math>, then <math>w(x)</math> is</p>
(A)	$e^{-3x}$
(B)	$e^{-2x}$
(C)	$e^{2x}$
(D)	$e^{3x}$



**GATE 2022 Mathematics (MA)**

Q.21	<p>The steady state solution for the heat equation</p> $\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} = 0, \quad 0 < x < 2, \quad t > 0,$ <p>with the initial condition <math>u(x, 0) = 0, \quad 0 &lt; x &lt; 2</math> and the boundary conditions <math>u(0, t) = 1</math> and <math>u(2, t) = 3, \quad t &gt; 0</math>, at <math>x = 1</math> is</p>
(A)	1
(B)	2
(C)	3
(D)	4



**GATE 2022 Mathematics (MA)**

<p>Q.22</p>	<p>Consider <math>([0, 1], T_1)</math>, where <math>T_1</math> is the subspace topology induced by the Euclidean topology on <math>\mathbb{R}</math>, and let <math>T_2</math> be <i>any</i> topology on <math>[0, 1]</math>. Consider the following statements:</p> <p><b>P</b> : If <math>T_1</math> is a proper subset of <math>T_2</math>, then <math>([0, 1], T_2)</math> is not compact.</p> <p><b>Q</b> : If <math>T_2</math> is a proper subset of <math>T_1</math>, then <math>([0, 1], T_2)</math> is not Hausdorff.</p> <p>Then</p>
<p>(A)</p>	<p><b>P</b> is TRUE and <b>Q</b> is FALSE</p>
<p>(B)</p>	<p>Both <b>P</b> and <b>Q</b> are TRUE</p>
<p>(C)</p>	<p>Both <b>P</b> and <b>Q</b> are FALSE</p>
<p>(D)</p>	<p><b>P</b> is FALSE and <b>Q</b> is TRUE</p>



**GATE 2022 Mathematics (MA)**

Q.23	Let $p : ([0, 1], T_1) \rightarrow (\{0, 1\}, T_2)$ be the quotient map, arising from the characteristic function on $[\frac{1}{2}, 1]$ , where $T_1$ is the subspace topology induced by the Euclidean topology on $\mathbb{R}$ . Which of the following statements is TRUE?
(A)	$p$ is an open map but not a closed map
(B)	$p$ is a closed map but not an open map
(C)	$p$ is a closed map as well as an open map
(D)	$p$ is neither an open map nor a closed map





**GATE 2022 Mathematics (MA)**

Q.24	<p>Set <math>X_n := \mathbb{R}</math> for each <math>n \in \mathbb{N}</math>. Define <math>Y := \prod_{n \in \mathbb{N}} X_n</math>. Endow <math>Y</math> with the product topology, where the topology on each <math>X_n</math> is the Euclidean topology. Consider the set</p> $\Delta = \{(x, x, x, \dots) \mid x \in \mathbb{R}\}$ <p>with the subspace topology induced from <math>Y</math>. Which of the following statements is TRUE?</p>
(A)	$\Delta$ is open in $Y$
(B)	$\Delta$ is locally compact
(C)	$\Delta$ is dense in $Y$
(D)	$\Delta$ is disconnected

**GATE 2022 Mathematics (MA)**

Q.25	<p>Consider the linear system of equations <math>Ax = b</math> with</p> $A = \begin{pmatrix} 3 & 1 & 1 \\ 1 & 4 & 1 \\ 2 & 0 & 3 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix}.$ <p>Which of the following statements are TRUE?</p>
(A)	<p>The Jacobi iterative matrix is <math>\begin{pmatrix} 0 &amp; 1/4 &amp; 1/3 \\ 1/3 &amp; 0 &amp; 1/3 \\ 2/3 &amp; 0 &amp; 0 \end{pmatrix}</math></p>
(B)	<p>The Jacobi iterative method converges for any initial vector</p>
(C)	<p>The Gauss-Seidel iterative method converges for any initial vector</p>
(D)	<p>The spectral radius of the Jacobi iterative matrix is less than 1</p>



GATE 2022 Mathematics (MA)

Q.26	The number of non-isomorphic abelian groups of order $2^2 \cdot 3^3 \cdot 5^4$ is _____.



GATE 2022 Mathematics (MA)

Q.27	The number of subgroups of a cyclic group of order 12 is _____.



GATE 2022 Mathematics (MA)

Q.28	The radius of convergence of the series $\sum_{n \geq 0} 3^{n+1} z^{2n}, z \in \mathbb{C}$ is _____ (round off to TWO decimal places).



GATE 2022 Mathematics (MA)

Q.29	The number of zeros of the polynomial $2z^7 - 7z^5 + 2z^3 - z + 1$ in the unit disc $\{z \in \mathbb{C} :  z  < 1\}$ is _____.



GATE 2022 Mathematics (MA)

Q.30	If $P(x)$ is a polynomial of degree 5 and $\alpha = \sum_{i=0}^6 P(x_i) \left( \prod_{j=0, j \neq i}^6 (x_i - x_j)^{-1} \right),$ where $x_0, x_1, \dots, x_6$ are distinct points in the interval $[2, 3]$ , then the value of $\alpha^2 - \alpha + 1$ is _____.



GATE 2022 Mathematics (MA)

Q.31	The maximum value of $f(x, y) = 49 - x^2 - y^2$ on the line $x + 3y = 10$ is _____.





GATE 2022 Mathematics (MA)

Q.32	If the function $f(x, y) = x^2 + xy + y^2 + \frac{1}{x} + \frac{1}{y}, x \neq 0, y \neq 0$ attains its local minimum value at the point $(a, b)$ , then the value of $a^3 + b^3$ is _____ (round off to TWO decimal places).



GATE 2022 Mathematics (MA)

<p>Q.33</p>	<p>If the ordinary differential equation</p> $x^2 \frac{d^2\phi}{dx^2} + x \frac{d\phi}{dx} + x^2\phi = 0, \quad x > 0$ <p>has a solution of the form <math>\phi(x) = x^r \sum_{n=0}^{\infty} a_n x^n</math>, where <math>a_n</math>'s are constants and <math>a_0 \neq 0</math>, then the value of <math>r^2 + 1</math> is _____.</p>



**GATE 2022 Mathematics (MA)**

Q.34	The Bessel functions $J_\alpha(x)$ , $x > 0$ , $\alpha \in \mathbb{R}$ satisfy $J_{\alpha-1}(x) + J_{\alpha+1}(x) = \frac{2\alpha}{x} J_\alpha(x)$ . Then, the value of $(\pi J_{\frac{3}{2}}(\pi))^2$ is _____.



GATE 2022 Mathematics (MA)

<p>Q.35</p>	<p>The partial differential equation</p> $7\frac{\partial^2 u}{\partial x^2} + 16\frac{\partial^2 u}{\partial x \partial y} + 4\frac{\partial^2 u}{\partial y^2} = 0$ <p>is transformed to</p> $A\frac{\partial^2 u}{\partial \xi^2} + B\frac{\partial^2 u}{\partial \xi \partial \eta} + C\frac{\partial^2 u}{\partial \eta^2} = 0,$ <p>using <math>\xi = y - 2x</math> and <math>\eta = 7y - 2x</math>. Then, the value of <math>\frac{1}{12^3}(B^2 - 4AC)</math> is _____.</p>

**GATE 2022 Mathematics (MA)**

**Q.36 – Q.65 Carry TWO marks each.**

Q.36	Let $\mathbb{R}[X]$ denote the ring of polynomials in $X$ with real coefficients. Then, the quotient ring $\mathbb{R}[X]/(X^4 + 4)$ is
(A)	a field
(B)	an integral domain, but not a field
(C)	not an integral domain, but has 0 as the only nilpotent element
(D)	a ring which contains non-zero nilpotent elements

GATE 2022 Mathematics (MA)

<p>Q.37</p>	<p>Consider the following conditions on two proper non-zero ideals <math>J_1</math> and <math>J_2</math> of a non-zero commutative ring <math>R</math>.</p> <p><b>P:</b> For any <math>r_1, r_2 \in R</math>, there exists a unique <math>r \in R</math> such that <math>r - r_1 \in J_1</math> and <math>r - r_2 \in J_2</math>.</p> <p><b>Q:</b> <math>J_1 + J_2 = R</math></p> <p>Then, which of the following statements is TRUE?</p>
<p>(A)</p>	<p><b>P</b> implies <b>Q</b> but <b>Q</b> does not imply <b>P</b></p>
<p>(B)</p>	<p><b>Q</b> implies <b>P</b> but <b>P</b> does not imply <b>Q</b></p>
<p>(C)</p>	<p><b>P</b> implies <b>Q</b> and <b>Q</b> implies <b>P</b></p>
<p>(D)</p>	<p><b>P</b> does not imply <b>Q</b> and <b>Q</b> does not imply <b>P</b></p>

**GATE 2022 Mathematics (MA)**

Q.38	Let $f : [-\pi, \pi] \rightarrow \mathbb{R}$ be a continuous function such that $f(x) > \frac{f(0)}{2}$ , $ x  < \delta$ for some $\delta$ satisfying $0 < \delta < \pi$ . Define $P_{n,\delta}(x) = (1 + \cos x - \cos \delta)^n$ , for $n = 1, 2, 3, \dots$ . Then, which of the following statements is TRUE?
(A)	$\lim_{n \rightarrow \infty} \int_0^{2\delta} f(x) P_{n,\delta}(x) dx = 0$
(B)	$\lim_{n \rightarrow \infty} \int_{-2\delta}^0 f(x) P_{n,\delta}(x) dx = 0$
(C)	$\lim_{n \rightarrow \infty} \int_{-\delta}^{\delta} f(x) P_{n,\delta}(x) dx = 0$
(D)	$\lim_{n \rightarrow \infty} \int_{[-\pi, \pi] \setminus [-\delta, \delta]} f(x) P_{n,\delta}(x) dx = 0$

GATE 2022 Mathematics (MA)

<p>Q.39</p>	<p><b>P</b> : Suppose that <math>\sum_{n=0}^{\infty} a_n x^n</math> converges at <math>x = -3</math> and diverges at <math>x = 6</math>. Then <math>\sum_{n=0}^{\infty} (-1)^n a_n</math> converges.</p> <p><b>Q</b>: The interval of convergence of the series <math>\sum_{n=2}^{\infty} \frac{(-1)^n x^n}{4^n \log_e n}</math> is <math>[-4, 4]</math>.</p> <p>Which of the following statements is TRUE?</p>
<p>(A)</p>	<p><b>P</b> is true and <b>Q</b> is true</p>
<p>(B)</p>	<p><b>P</b> is false and <b>Q</b> is false</p>
<p>(C)</p>	<p><b>P</b> is true and <b>Q</b> is false</p>
<p>(D)</p>	<p><b>P</b> is false and <b>Q</b> is true</p>





GATE 2022 Mathematics (MA)

Q.40	<p>Let</p> $f_n(x) = \frac{x^2}{x^2 + (1 - nx)^2}, \quad x \in [0, 1], \quad n = 1, 2, 3, \dots$ <p>Then, which of the following statements is TRUE?</p>
(A)	$\{f_n\}$ is not equicontinuous on $[0, 1]$
(B)	$\{f_n\}$ is uniformly convergent on $[0, 1]$
(C)	$\{f_n\}$ is equicontinuous on $[0, 1]$
(D)	$\{f_n\}$ is uniformly bounded and has a subsequence converging uniformly on $[0, 1]$



GATE 2022 Mathematics (MA)

Q.41	Let $(\mathbb{Q}, d)$ be the metric space with $d(x, y) =  x - y $ . Let $E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}$ . Then, the set $E$ is
(A)	closed but not compact
(B)	not closed but compact
(C)	compact
(D)	neither closed nor compact



**GATE 2022 Mathematics (MA)**

Q.42	Let $T : L^2[-1, 1] \rightarrow L^2[-1, 1]$ be defined by $Tf = \tilde{f}$ , where $\tilde{f}(x) = f(-x)$ almost everywhere. If $M$ is the kernel of $I - T$ , then the distance between the function $\phi(t) = e^t$ and $M$ is
(A)	$\frac{1}{2}\sqrt{(e^2 - e^{-2} + 4)}$
(B)	$\frac{1}{2}\sqrt{(e^2 - e^{-2} - 2)}$
(C)	$\frac{1}{2}\sqrt{(e^2 - 4)}$
(D)	$\frac{1}{2}\sqrt{(e^2 - e^{-2} - 4)}$



**GATE 2022 Mathematics (MA)**

Q.43	Let $X$ , $Y$ and $Z$ be Banach spaces. Suppose that $T : X \rightarrow Y$ is linear and $S : Y \rightarrow Z$ is linear, bounded and injective. In addition, if $S \circ T : X \rightarrow Z$ is bounded, then, which of the following statements is TRUE?
(A)	$T$ is surjective
(B)	$T$ is bounded but not continuous
(C)	$T$ is bounded
(D)	$T$ is not bounded

GATE 2022 Mathematics (MA)

<p>Q.44</p>	<p>The first derivative of a function <math>f \in C^\infty(-3, 3)</math> is approximated by an interpolating polynomial of degree 2, using the data</p> $(-1, f(-1)), (0, f(0)) \text{ and } (2, f(2)).$ <p>It is found that</p> $f'(0) \approx -\frac{2}{3}f(-1) + \alpha f(0) + \beta f(2).$ <p>Then, the value of <math>\frac{1}{\alpha\beta}</math> is</p>
<p>(A)</p>	<p>3</p>
<p>(B)</p>	<p>6</p>
<p>(C)</p>	<p>9</p>
<p>(D)</p>	<p>12</p>



**GATE 2022 Mathematics (MA)**

Q.45	The work done by the force $F = (x + y)\hat{i} - (x^2 + y^2)\hat{j}$ , where $\hat{i}$ and $\hat{j}$ are unit vectors in $\overrightarrow{OX}$ and $\overrightarrow{OY}$ directions, respectively, along the upper half of the circle $x^2 + y^2 = 1$ from $(1, 0)$ to $(-1, 0)$ in the $xy$ -plane is
(A)	$-\pi$
(B)	$-\frac{\pi}{2}$
(C)	$\frac{\pi}{2}$
(D)	$\pi$

**GATE 2022 Mathematics (MA)**

<p>Q.46</p>	<p>Let <math>u(x, t)</math> be the solution of the wave equation</p> $\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} = 0, \quad 0 < x < \pi, \quad t > 0,$ <p>with the initial conditions</p> $u(x, 0) = \sin x + \sin 2x + \sin 3x, \quad \frac{\partial u}{\partial t}(x, 0) = 0, \quad 0 < x < \pi$ <p>and the boundary conditions <math>u(0, t) = u(\pi, t) = 0, \quad t \geq 0</math>. Then, the value of <math>u\left(\frac{\pi}{2}, \pi\right)</math> is</p>
<p>(A)</p>	<p><math>-1/2</math></p>
<p>(B)</p>	<p><math>0</math></p>
<p>(C)</p>	<p><math>1/2</math></p>
<p>(D)</p>	<p><math>1</math></p>



**GATE 2022 Mathematics (MA)**

Q.47	<p>Let <math>T : \mathbb{R}^2 \rightarrow \mathbb{R}^2</math> be a linear transformation defined by</p> $T((1, 2)) = (1, 0) \quad \text{and} \quad T((2, 1)) = (1, 1).$ <p>For <math>p, q \in \mathbb{R}</math>, let <math>T^{-1}((p, q)) = (x, y)</math>.</p> <p>Which of the following statements is TRUE?</p>
(A)	$x = p - q; y = 2p - q$
(B)	$x = p + q; y = 2p - q$
(C)	$x = p + q; y = 2p + q$
(D)	$x = p - q; y = 2p + q$



**GATE 2022 Mathematics (MA)**

<p>Q.48</p>	<p>Let <math>y = (\alpha, -1)^T</math>, <math>\alpha \in \mathbb{R}</math> be a feasible solution for the dual problem of the linear programming problem</p> $\begin{aligned} &\text{Maximize:} && 5x_1 + 12x_2 \\ &\text{subject to:} && x_1 + 2x_2 + x_3 \leq 10 \\ &&& 2x_1 - x_2 + 3x_3 = 8 \\ &&& x_1, x_2, x_3 \geq 0. \end{aligned}$ <p>Which of the following statements is TRUE?</p>
<p>(A)</p>	<p><math>\alpha &lt; 3</math></p>
<p>(B)</p>	<p><math>3 \leq \alpha &lt; 5.5</math></p>
<p>(C)</p>	<p><math>5.5 \leq \alpha &lt; 7</math></p>
<p>(D)</p>	<p><math>\alpha \geq 7</math></p>



GATE 2022 Mathematics (MA)

Q.49	Let $K$ denote the subset of $\mathbb{C}$ consisting of elements algebraic over $\mathbb{Q}$ . Then, which of the following statements are TRUE?
(A)	No element of $\mathbb{C} \setminus K$ is algebraic over $\mathbb{Q}$
(B)	$K$ is an algebraically closed field
(C)	For any bijective ring homomorphism $f : \mathbb{C} \rightarrow \mathbb{C}$ , we have $f(K) = K$
(D)	There is no bijection between $K$ and $\mathbb{Q}$



**GATE 2022 Mathematics (MA)**

Q.50	Let $T$ be a Möbius transformation such that $T(0) = \alpha, T(\alpha) = 0$ and $T(\infty) = -\alpha$ , where $\alpha = (-1 + i)/\sqrt{2}$ . Let $L$ denote the straight line passing through the origin with slope $-1$ , and let $C$ denote the circle of unit radius centred at the origin. Then, which of the following statements are TRUE?
(A)	$T$ maps $L$ to a straight line
(B)	$T$ maps $L$ to a circle
(C)	$T^{-1}$ maps $C$ to a straight line
(D)	$T^{-1}$ maps $C$ to a circle



**GATE 2022 Mathematics (MA)**

Q.51	Let $a > 0$ . Define $D_a : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ by $(D_a f)(x) = \frac{1}{\sqrt{a}} f\left(\frac{x}{a}\right)$ , almost everywhere, for $f \in L^2(\mathbb{R})$ . Then, which of the following statements are TRUE?
(A)	$D_a$ is a linear isometry
(B)	$D_a$ is a bijection
(C)	$D_a \circ D_b = D_{a+b}$ , $b > 0$
(D)	$D_a$ is bounded from below



**GATE 2022 Mathematics (MA)**

Q.52	Let $\{\phi_0, \phi_1, \phi_2, \dots\}$ be an orthonormal set in $L^2[-1, 1]$ such that $\phi_n = C_n P_n$ , where $C_n$ is a constant and $P_n$ is the Legendre polynomial of degree $n$ , for each $n \in \mathbb{N} \cup \{0\}$ . Then, which of the following statements are TRUE?
(A)	$\phi_6(1) = 1$
(B)	$\phi_7(-1) = 1$
(C)	$\phi_7(1) = \sqrt{\frac{15}{2}}$
(D)	$\phi_6(-1) = \sqrt{\frac{13}{2}}$



GATE 2022 Mathematics (MA)

Q.53	Let $X = (\mathbb{R}, T)$ , where $T$ is the smallest topology on $\mathbb{R}$ in which all the singleton sets are closed. Then, which of the following statements are TRUE?
(A)	$[0, 1)$ is compact in $X$
(B)	$X$ is not first countable
(C)	$X$ is second countable
(D)	$X$ is first countable



**GATE 2022 Mathematics (MA)**

Q.54	<p>Consider <math>(\mathbb{Z}, T)</math>, where <math>T</math> is the topology generated by sets of the form</p> $A_{m,n} = \{m + nk \mid k \in \mathbb{Z}\},$ <p>for <math>m, n \in \mathbb{Z}</math> and <math>n \neq 0</math>. Then, which of the following statements are TRUE?</p>
(A)	$(\mathbb{Z}, T)$ is connected
(B)	Each $A_{m,n}$ is a closed subset of $(\mathbb{Z}, T)$
(C)	$(\mathbb{Z}, T)$ is Hausdorff
(D)	$(\mathbb{Z}, T)$ is metrizable

**GATE 2022 Mathematics (MA)**

<p>Q.55</p>	<p>Let <math>A \in \mathbb{R}^{m \times n}</math>, <math>c \in \mathbb{R}^n</math> and <math>b \in \mathbb{R}^m</math>. Consider the linear programming primal problem</p> $\begin{aligned} &\text{Minimize: } c^T x \\ &\text{subject to: } Ax = b \\ &\quad \quad \quad x \geq 0. \end{aligned}$ <p>Let <math>x^0</math> and <math>y^0</math> be feasible solutions of the primal and its dual, respectively. Which of the following statements are TRUE?</p>
<p>(A)</p>	<p><math>c^T x^0 \geq b^T y^0</math></p>
<p>(B)</p>	<p><math>c^T x^0 = b^T y^0</math></p>
<p>(C)</p>	<p>If <math>c^T x^0 = b^T y^0</math>, then <math>x^0</math> is optimal for the primal</p>
<p>(D)</p>	<p>If <math>c^T x^0 = b^T y^0</math>, then <math>y^0</math> is optimal for the dual</p>



**GATE 2022 Mathematics (MA)**

<p>Q.56</p>	<p>Consider <math>\mathbb{R}^3</math> as a vector space with the usual operations of vector addition and scalar multiplication. Let <math>x \in \mathbb{R}^3</math> be denoted by <math>x = (x_1, x_2, x_3)</math>. Define subspaces <math>W_1</math> and <math>W_2</math> by</p> $W_1 := \{x \in \mathbb{R}^3 : x_1 + 2x_2 - x_3 = 0\}$ <p>and</p> $W_2 := \{x \in \mathbb{R}^3 : 2x_1 + 3x_3 = 0\}.$ <p>Let <math>\dim(U)</math> denote the dimension of the subspace <math>U</math>.</p> <p>Which of the following statements are TRUE?</p>
<p>(A)</p>	<p><math>\dim(W_1) = \dim(W_2)</math></p>
<p>(B)</p>	<p><math>\dim(W_1) + \dim(W_2) - \dim(\mathbb{R}^3) = 1</math></p>
<p>(C)</p>	<p><math>\dim(W_1 + W_2) = 2</math></p>
<p>(D)</p>	<p><math>\dim(W_1 \cap W_2) = 1</math></p>



GATE 2022 Mathematics (MA)

<p>Q.57</p>	<p>Three companies <math>C_1, C_2</math> and <math>C_3</math> submit bids for three jobs <math>J_1, J_2</math> and <math>J_3</math>. The costs involved per unit are given in the table below:</p> <table style="margin-left: auto; margin-right: auto;"> <tr> <td></td> <td style="text-align: center;"><math>J_1</math></td> <td style="text-align: center;"><math>J_2</math></td> <td style="text-align: center;"><math>J_3</math></td> </tr> <tr> <td style="text-align: center;"><math>C_1</math></td> <td style="border: 1px solid black; text-align: center;">10</td> <td style="border: 1px solid black; text-align: center;">12</td> <td style="border: 1px solid black; text-align: center;">8</td> </tr> <tr> <td style="text-align: center;"><math>C_2</math></td> <td style="border: 1px solid black; text-align: center;">9</td> <td style="border: 1px solid black; text-align: center;">15</td> <td style="border: 1px solid black; text-align: center;">10</td> </tr> <tr> <td style="text-align: center;"><math>C_3</math></td> <td style="border: 1px solid black; text-align: center;">15</td> <td style="border: 1px solid black; text-align: center;">10</td> <td style="border: 1px solid black; text-align: center;">9</td> </tr> </table> <p>Then, the cost of the optimal assignment is _____.</p>		$J_1$	$J_2$	$J_3$	$C_1$	10	12	8	$C_2$	9	15	10	$C_3$	15	10	9
	$J_1$	$J_2$	$J_3$														
$C_1$	10	12	8														
$C_2$	9	15	10														
$C_3$	15	10	9														

GATE 2022 Mathematics (MA)

<p>Q.58</p>	<p>The initial value problem <math>\frac{dy}{dx} = f(x, y)</math>, <math>y(x_0) = y_0</math> is solved by using the following second order Runge-Kutta method:</p> $K_1 = hf(x_i, y_i)$ $K_2 = hf(x_i + \alpha h, y_i + \beta K_1)$ $y_{i+1} = y_i + \frac{1}{4}(K_1 + 3K_2), \quad i \geq 0,$ <p>where <math>h</math> is the uniform step length between the points <math>x_0, x_1, \dots, x_n</math> and <math>y_i = y(x_i)</math>. The value of the product <math>\alpha\beta</math> is _____ (round off to TWO decimal places).</p>



**GATE 2022 Mathematics (MA)**

Q.59	The surface area of the paraboloid $z = x^2 + y^2$ between the planes $z = 0$ and $z = 1$ is _____ (round off to ONE decimal place).



GATE 2022 Mathematics (MA)

Q.60	The rate of change of $f(x, y, z) = x + x \cos z - y \sin z + y$ at $P_0$ in the direction from $P_0(2, -1, 0)$ to $P_1(0, 1, 2)$ is _____.



GATE 2022 Mathematics (MA)

<p>Q.61</p>	<p>If the Laplace equation</p> $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \quad 1 < x < 2, \quad 1 < y < 2$ <p>with the boundary conditions</p> $\frac{\partial u}{\partial x}(1, y) = y, \quad \frac{\partial u}{\partial x}(2, y) = 5, \quad 1 < y < 2$ <p>and</p> $\frac{\partial u}{\partial y}(x, 1) = \frac{\alpha x^2}{7}, \quad \frac{\partial u}{\partial y}(x, 2) = x, \quad 1 < x < 2$ <p>has a solution, then the constant <math>\alpha</math> is _____.</p>



GATE 2022 Mathematics (MA)

Q.62	Let $u(x, y)$ be the solution of the first order partial differential equation $x \frac{\partial u}{\partial x} + (x^2 + y) \frac{\partial u}{\partial y} = u, \text{ for all } x, y \in \mathbb{R}$ satisfying $u(2, y) = y - 4, y \in \mathbb{R}$ . Then, the value of $u(1, 2)$ is _____.



GATE 2022 Mathematics (MA)

Q.63	The optimal value for the linear programming problem $\text{Maximize: } 6x_1 + 5x_2$ $\text{subject to: } 3x_1 + 2x_2 \leq 12$ $-x_1 + x_2 \leq 1$ $x_1, x_2 \geq 0$ is _____.



**GATE 2022 Mathematics (MA)**

Q.64	<p>A certain product is manufactured by plants <math>P_1, P_2</math> and <math>P_3</math> whose capacities are 15, 25 and 10 units, respectively. The product is shipped to markets <math>M_1, M_2, M_3</math> and <math>M_4</math>, whose requirements are 10, 10, 10 and 20, respectively. The transportation costs per unit are given in the table below.</p> <table style="margin: 20px auto; border-collapse: collapse;"> <tr> <td style="border: none;"></td> <td style="border: 1px solid black; padding: 5px;"><math>M_1</math></td> <td style="border: 1px solid black; padding: 5px;"><math>M_2</math></td> <td style="border: 1px solid black; padding: 5px;"><math>M_3</math></td> <td style="border: 1px solid black; padding: 5px;"><math>M_4</math></td> <td style="border: none;"></td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;"><math>P_1</math></td> <td style="border: 1px solid black; padding: 5px; text-align: center;">1</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">3</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">1</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">3</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">15</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;"><math>P_2</math></td> <td style="border: 1px solid black; padding: 5px; text-align: center;">2</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">2</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">4</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">1</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">25</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;"><math>P_3</math></td> <td style="border: 1px solid black; padding: 5px; text-align: center;">2</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">1</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">1</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">2</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">10</td> </tr> <tr> <td style="border: none;"></td> <td style="border: 1px solid black; padding: 5px; text-align: center;">10</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">10</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">10</td> <td style="border: 1px solid black; padding: 5px; text-align: center;">20</td> <td style="border: none;"></td> </tr> </table> <p>Then the cost corresponding to the starting basic solution by the Northwest-corner method is _____.</p>		$M_1$	$M_2$	$M_3$	$M_4$		$P_1$	1	3	1	3	15	$P_2$	2	2	4	1	25	$P_3$	2	1	1	2	10		10	10	10	20	
	$M_1$	$M_2$	$M_3$	$M_4$																											
$P_1$	1	3	1	3	15																										
$P_2$	2	2	4	1	25																										
$P_3$	2	1	1	2	10																										
	10	10	10	20																											



**GATE 2022 Mathematics (MA)**

Q.65	Let $M$ be a $3 \times 3$ real matrix such that $M^2 = 2M + 3I$ . If the determinant of $M$ is $-9$ , then the trace of $M$ equals _____.