


Titration Curve of Weak Acid with Strong Base



Titration Curve of Weak Acid with Strong Base


In the titration of weak acid versus strong base, protons get transferred from weak acid to strong base takes place. Let us take an example of titration of acetic acid (weak acid) with sodium hydroxide (strong base).



The titration curve generally reflects the strength of the base and acid in which a change in pH takes place. For acetic acid and sodium hydroxide, it is observed that:

- I. In the beginning, there will be a sharp increase in pH because of the common ion effect.
- II. After this, a change in curve takes place very slowly and gradually because the solution will act as a buffer now.
- III. At the middle of the curve, half neutralization will take place at which the concentration of both weak acid and its conjugate base becomes equal.
- IV. At the equivalence point, $\text{pH} > 7$ because after the addition of NaOH, all acid gets converted to its conjugate base and at this point due to the common ion effect, the reaction will move backward and produce hydroxide ion.
- V. After the equivalence point, the curve becomes typical.


Graphical representation:




Buffer solution - A mixture of a weak acid along with its conjugate base or vice versa is known as a buffer solution. With the addition of a small amount of either acid or base, the buffer solution resists a change in pH.

Working on a buffer solution

A mixture of acetic acid and sodium hydroxide acts as a buffer solution. When a small amount of either of these takes place, the concentration of hydronium ion remains constant. In addition to a small amount of NaOH, $[\text{OH}]^-$ will react with a few $[\text{H}]^+$ ions. As a result, more of the acetic acid tends to react with water due to which $[\text{H}]^+$ ions remain restored to their original value.



On the addition of HCl acid, H^+ ions will react with acetate ions and this reaction can be represented as:



Again, there is a very minute change in $[\text{H}]^+$ ions due to which pH remains the same. A mixture of acetic acid and sodium hydroxide is an acidic buffer solution (weak acid and strong base). The pH for such solutions can be calculated by using Henderson- Hasselbalch equation.

$$\text{pH} = \text{pKa} + \log \frac{[\text{conjugate base or salt}]}{[\text{acid}]}$$

CSIR NET

Life Science 2022

A Foundation Course

Complete Prep of Life Science for June 2022 Aspirants

Why take this course?

- 400+ Hrs Live Classes & Doubt Sessions for complete conceptual clarity
- 3000+ Practice Questions covering all levels of difficulty
- 20+ Unit Wise Study Modules & Mind Maps
- 50+ Full Mock Tests, Chapter Wise Mock Tests, PYQs Mock Tests

